

## MEETING ABSTRACT

## Open Access

# Pivotal vascular homeostatic role for endothelium-derived C-type natriuretic peptide (CNP)

Adrian J Hobbs

From 7th International Conference on cGMP Generators, Effectors and Therapeutic Implications  
Trier, Germany. 19-21 June 2015

## Background

The endothelium plays a fundamental role in maintaining vascular homeostasis by releasing factors that regulate local blood flow, systemic blood pressure, and the reactivity of leukocytes and platelets. Accordingly, endothelial dysfunction underpins many cardiovascular diseases, including hypertension, myocardial infarction and stroke. C-type natriuretic peptide (CNP) is a paracrine mediator that possesses a unique vaso- and cardio-protective pharmacodynamic profile; however, there is a paucity of information regarding a physiological role for CNP within the cardiovascular system.

## Methods and principal results

Herein, we generated a novel endothelial cell-specific CNP knockout (ecCNP KO) mouse to investigate a potential vasoprotective role of endogenous CNP in vivo. Mice lacking endothelial CNP exhibited a fundamental, multi-faceted vascular dysfunction, including impaired endothelium-dependent dilatation, hypertension, increased leukocyte & platelet reactivity, accelerated atherogenesis and aneurysm. Moreover, the aberrant vascular phenotype observed in ecCNP KO animals was more closely recapitulated in mice with global deletion of natriuretic peptide receptor (NPR)-C than animals lacking guanylate cyclase-coupled NPR-B. In accord, we designed & developed a series of novel, small molecule NPR-C agonists, which are vasorelaxant in vitro and reduce blood pressure in vivo.

## Conclusion

These data identify a mechanism explaining genome-wide association studies linking the NPR-C (Npr3) and Furin (a pro-protein convertase obligatory in the bioactivation of CNP) loci with hypertension, and establish the importance of CNP-NPR-C signalling in preserving vascular homeostasis and as a disease-modifying drug target.

This work was supported by the Wellcome Trust and UCL Business PLC.

Published: 2 September 2015

doi:10.1186/2050-6511-16-S1-A27

**Cite this article as:** Hobbs: Pivotal vascular homeostatic role for endothelium-derived C-type natriuretic peptide (CNP). *BMC Pharmacology and Toxicology* 2015 **16**(Suppl 1):A27.

## Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)



Correspondence: [a.j.hobbs@qmul.ac.uk](mailto:a.j.hobbs@qmul.ac.uk)  
William Harvey Research Institute, London, UK



© 2015 Hobbs This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated.